

Date of Procedure: _____ Time: _____

(IMPORTANT: You must arrive 30 minutes prior to your appointment and **arrange for a friend or relative to drive you to and from your procedure.**)

Phone: (847)247-0187

Arrive at: _____

Location: **Gurnee Endoscopy**
731 S. IL Route 21, Suite 150
Gurnee, IL 60031

You have been scheduled for a colonoscopy. A colonoscopy is an examination of the colon (large bowel) with a lighted flexible tube called a colonoscope. Your efforts at cleansing your colon are essential for an accurate examination.

Colonoscopy Preparation with SUTAB:

Please purchase the items 5 days before your procedure:

2 Bisacodyl 5mg Tablets (Laxatives) Dulcolax

1 SUTAB Preparation Kit (By prescription from pharmacy)

(DO NOT FOLLOW THE INSTRUCTIONS ON THE BOX! PLEASE FOLLOW THESE INSTRUCTIONS!)

Read these instructions carefully at least one week prior to your exam. For more information about the benefits and risks of colonoscopy, please refer to the information pamphlet given to you, or call our office for additional information.

Please Contact your PCP/Cardiologist for cardiac clearance if any of the following relate to your current health:

- You are currently being treated for a cardiac condition.
- You have a **cardiac stent** and are taking a prescription medication.
- If you are currently taking any "blood thinners"; such as Plavix, Coumadin or Lovenox to receive medication.
- You have a pacemaker or defibrillator please let our office know immediately.

Medication and Diet Instructions:

Five days before the procedure:

1. Stop taking medications listed below:
 - Coumadin (Warfarin) Effient (prasugrel)
 - Plavix (clopidogrel) Brilinta (ticagrelor)
 - Pradaxa (dabigatran) - 2 Days Xarelto (Rivaroxaban) - 2 Days
 - Lovenox - 1 Day Eliquis (Apixaban) - 2 Days
 - Heparin
2. If you have an artificial heart valve, you need to talk to your primary care physician or cardiologist about using an alternative anticoagulant. Make sure the doctor doing your colonoscopy knows that you have a prosthetic valve.

Cardiac Patients

If you are being treated for hypertension/high blood pressure, continue to take your medication as directed. On the day of the procedure take the medication with **ONLY** a sip of water.

Diabetic Patients

If you are taking an oral hypoglycemic ("sugar pill"), **do not take** it the day of your procedure. If you are taking insulin, please take 1/2 of your prescribed a.m. insulin dose, the day before your procedure. Do NOT take insulin the day of. Call your PCP if you have any questions about your medicine during your preparation period.

General Information

1. You must arrive between **30 mins** before the scheduled appointment according to the guidelines of the procedure location. Your entire stay for the procedure will be approximately 1 to 1 ½ hours.
 2. You are NOT allowed to take a taxi or public transportation home after the procedure. A friend or relative will need to drive you home, and someone will need to be with you for at least two hours to make sure there are no problems from the sedation or the procedure.
 3. After the procedure, do not drive a car for the remainder of the day and do not make any important decisions until the following day.
 4. Biopsy results are usually available within five business days.
1. **4Patients with pacemakers** need to inform our office when booking this appointment.
 2. **Patients with internal defibrillators** must have the procedure at Condell Medical

***** Three days before the procedure: Avoid eating nuts, seeds, popcorn, grapes, green peas, beans, and tomatoes. *******

Two Days before your procedure:

_____ Take 2 Dulcolax tablets at bedtime.

Day before the procedure: _____

Start drinking clear liquids in the morning and continue throughout the day. Clear liquids include black coffee/tea (WITH NO MILK). Soda, Gatorade, Propel, water, clear juices (apple juice, white grape, white cranberry), clear broth, popsicles, and jell-o (NO RED, BLUE OR PURPLE). **Do not eat any solid food while on this diet. Do not drink alcohol. No dairy products.** Deviating from this diet may prevent adequate preparation for the colon exam.

Take Dose 1 at 1 PM:

- Open 1 bottle of **12 tablets**. Fill the provided container with **16 ounces of water** (up to the fill line). Swallow each tablet with a sip of water and drink the entire amount of water **over 15 to 20 minutes**.
- Approximately **1 hour after** the last tablet is ingested, fill the provided container again with **16 ounces of any clear liquid** (up to the fill line), and drink the entire amount over 30 minutes.
- Approximately **30 minutes after** finishing the second container of clear liquid, fill the provided container with **16 ounces of any clear liquid** (up to the fill line), and drink the entire amount over 30 minutes.
- **Two hours later drink another 16 ounces of any clear liquid!**

Take Dose 2 at 8 PM:

- Open 1 bottle of **12 tablets**. Fill the provided container with **16 ounces of water** (up to the fill line). Swallow each tablet with a sip of water and drink the entire amount of water **over 15 to 20 minutes**.
- Approximately **1 hour after** the last tablet is ingested, fill the provided container again with **16 ounces of any clear liquid** (up to the fill line), and drink the entire amount over 30 minutes.
- Approximately **30 minutes after** finishing the second container of clear liquid, fill the provided container with **16 ounces of any clear liquid** (up to the fill line), and drink the entire amount over 30 minutes.
- **Two hours later drink another 16 ounces of any clear liquid!**

***ABSOLUTELY NOTHING BY MOUTH AFTER MIDNIGHT ***

